

Einstein-Podolsky-Rosen Paradox and Antiparticle

Guang-jiong Ni*

Department of Physics, Fudan University, Shanghai, 200433, P. R. China

Hong Guan†

Department of Physics, Zhongshan University, Guangzhou, 510275, P. R. China

The original version of Einstein-Podolsky-Rosen (EPR) paradox is discussed to show the completeness of Quantum Mechanics (QM). The unique solution leads to the wave function of antiparticle unambiguously, which implies the essential conformity between QM and Special Relativity (SR).

PACS: 03.65.Bz

Einstein, Podolsky and Rosen (EPR) in their famous paper in 1935 [1] insisted on querying that “Is the description of physical reality in quantum mechanics (QM) complete?” What EPR were discussing is a system composing of two spinless particles. It was a very strange example [2], see below. So quite naturally, beginning from Bohm [3], physicists have been turning to other experiments involving particle spin or photons. The investigations on these EPR experiments together with the analysis of Bell’s inequality [4] have been revealing that the prediction of QM is correct while the existence of Local Hidden Variable (LHV) is incompatible with the outcome of experiments. A recent experiment [5] even showed that the quantum correlation in an entangled state of two-photon system can be maintained over long distance exceeding 10 km. However, it seems to us that the original version of EPR paradox still remains to be answered.

The question EPR raised is as follows. As is well known in QM, the position of a particle, x , and its momentum (in one dimensional space) operator

$$\hat{p} = -i\hbar \frac{\partial}{\partial x} \quad (1)$$

have a commutation relation:

$$[x, \hat{p}] = i\hbar \quad (2)$$

Consider a two-particle system. Then the operators $(x_1 - x_2)$ and $(\hat{p}_1 + \hat{p}_2)$ will commute:

$$[x_1 - x_2, \hat{p}_1 + \hat{p}_2] = 0 \quad (3)$$

which means that they have a common eigenstate with eigenvalues

$$x_1 - x_2 = a = \text{const} \quad (4)$$

and

$$p_1 + p_2 = 0, \quad p_2 = -p_1 \quad (5)$$

Now question arises: What strange the state is!? Two particles are moving in opposite momentum directions while keeping their distance unchanged. Incredible! As stressed in Ref. [2]: “No one can figure out how to realize it”. “So there is no surprise that many people regarded the debate (between EPR and Bohr) as an idle talk leading only to empty result”. However, as we believe in the basic principles of QM, any difficulty must have its root. Is this implying something overlooked or unfinished in QM?

In this letter, we wish to propose an answer to this EPR paradox, which essentially prescribes the necessity of introducing the antiparticle state in QM and even dictates the form of its Wave Function (WF) unambiguously.

If the WF of particle 1 is written as usual:

$$\psi(x_1, t) = \exp\left\{\frac{i}{\hbar}(p_1 x_1 - E_1 t)\right\} \quad (6)$$

with momentum $p_1 > 0$ and energy $E_1 > 0$, then the particle 2 must be an antiparticle described by the following WF:

*E-mail: gjni@fudan.ac.cn

†E-mail: stdp@zsu.edu.cn

$$\psi_c(x_2, t) = \exp\left\{\frac{i}{\hbar}(p_2 x_2 - E_2 t)\right\} = \exp\left\{-\frac{i}{\hbar}(p_1 x_2 - E_1 t)\right\} \quad (7)$$

with $p_2 = -p_1 < 0$, $E_2 = -E_1 < 0$. But as proposed by Schwinger ([6], see also [7]) and stressed by one of the present authors, Ni ([8], [9]), we should view the WF of “negative energy state” of particle directly as the WF of its antiparticle with the corresponding antiparticle operators:

$$\hat{p}_c = i\hbar \frac{\partial}{\partial x}, \quad \hat{E}_c = -i\hbar \frac{\partial}{\partial t} \quad (8)$$

So the observed momentum and energy of antiparticle in state (7) are p_1 and E_1 respectively, precisely the same as that of the particle state (6). Therefore, the EPR state $\Psi = \psi\psi_c$ is composed of a particle-antiparticle pair with parallel momentum (p_1) and invariant distance ($x_1 - x_2$) between them. Now every thing is reasonable.

In most text books on QM, there is nearly no clear explanation for WF of antiparticle. Some times (and often implicitly), it is said that the WF of antiparticle with momentum p_1 is the same as that of particle, i.e., Eq. (6), one has to distinguish a positron from electron by adding some words. Alternatively, one often said that in the vacuum all the negative energy states of electron are filled. Once a “hole” is created in the “sea”, it would correspond to a positron.

In accompanying with the historic discovery of parity (P) violation by Lee-Yang [10] and Wu et. al. [11] in 1956, it was verified that the charge conjugate transformation (C) which brings an electron (with charge $-e$) to a positron (with charge e) is also violated [12]. In 1964, the combined CP violation was discovered in $K^0 - \bar{K}^0$ system whereas the CPT theorem remains valid. The recent experiment by CPLEAR Collaboration at CERN provides further direct observation of time-reversal (T) non-invariance in the neutral-kaon system [13] (see also [14]). What do all the above discoveries mean?

In our point of view, all the above discoveries imply that the individual definition of P, C or T inversion ceases to be meaningful as an observable transformation in physics. Of course, their definitions are still clearcut in mathematics. Let us perform the combined CPT transformation on the WF of a particle, either with or without spin. (see, e.g. in [15]). Because the C or T transformation contains a complex conjugation respectively, they cancel each other in the CPT transformation. Then what we obtain is essentially a transformation ($\vec{x} \rightarrow -\vec{x}, t \rightarrow -t$) while at the same time we should look at the WF describing an antiparticle. This is nothing but a statement of transforming Eq. (6) to Eq. (7). And this is why physicists have been gradually accepting the relation of a particle $|a\rangle$ with respect to its antiparticle $|\bar{a}\rangle$ being [16]

$$|\bar{a}\rangle = CPT|a\rangle \quad (9)$$

The reason why the observation Eqs. (6)-(8) had not attracted enough attention in physics community for so long a time is of two fold. First, the inertia of concept in physics proves its strength once again as witnessed in the history of science from time to time. Second, the importance of the above observation has not fully explored until its evolving into the following postulate: “The space-time reversal ($\vec{x} \rightarrow -\vec{x}, t \rightarrow -t$) transformation is equivalent to the transformation between particle and antiparticle.” This is a basic symmetry which should be respected in constructing the theory for all particles and fields. Thus a particle is always not pure. A particle state $\theta(\vec{x}, t)$ is always accompanied by its antiparticle state $\chi(\vec{x}, t)$. They have the simple relation:

$$\theta(-\vec{x}, -t) = \chi(\vec{x}, t) \quad (10)$$

The coupled equations should be invariant with respect to the transformation ($\vec{x} \rightarrow -\vec{x}, t \rightarrow -t$) together with Eq. (10).

Based on this symmetry, the essence of special relativity (SR) is explored and the one-body relativistic equation, i.e., the Dirac equation and Klein-Gordon equation, are derived [9]. Moreover, even to our surprise, we began to realize that the Stationary Schrödinger Equation (SSE) for many-body system is essentially relativistic as long as the eigenvalue in SSE

$$H\psi = \varepsilon\psi \quad (11)$$

(say for two-body case, $H = p^2/2\mu + V(r)$ with μ being the reduced mass) is related to the binding energy B ($B \equiv M - E$ with M and E being the rest mass and energy of whole system, $c = 1$) as follows ([17], [18]):

$$\varepsilon \equiv \frac{E^2 - M^2}{2M}, \quad B = M[1 - (1 + \frac{2\varepsilon}{M})^{1/2}] \quad (12)$$

It is interesting to see that the relativistic correction in Eq. (12) for Hydrogenlike atom is a small downward energy shift

$$\Delta E^{rel} = -\frac{\varepsilon^2}{2M}, \quad (\varepsilon = -\frac{Z^2\alpha^2\mu}{2n^2}) \quad (13)$$

which only accounts for $(-23.814MHz)$, or (-0.3%) in the magnitude of the so-called absolute Lamb shift (upward) $8172.86MHz$ of 1S state in Hydrogen atom [19]. Based on the new understanding of SSE, Eq (11), a simple noncovariant calculation on Lamb shift is performed in Ref [20] with high accuracy ($< 0.1\%$).

In summary, some discussions are in order:

(a) The original version of EPR paradox was a simple but acute problem querying on whether the QM is complete. If instead of Eq. (3) we consider

$$[x_1 + x_2, \hat{p}_1 - \hat{p}_2] = 0 \quad (14)$$

Then the correct answer turns out to be a particle and its antiparticle (c) moving in opposite direction with momentum p_1 and $p_2^{(c)} = -p_1$ and position x_1 and $x_2 = -x_1$. Such kind of experiments have been performed for many times. For example, the e^+ and e^- pair can be created by a high energy photon in the vicinity of heavy nuclei. The 0^+ excited state of ^{16}O nuclei lies higher $6MeV$ than the ground 0^+ state, so it can also give rise to pair creation of e^+ and e^- . Recently, the experiment at CERN again demonstrated the quantum correlation at a distance for kaon and antikaon system. They are entwined. Only until the measurement on p_1 direction to see a particle (or antiparticle), can one predict with 100 percent probability that an antiparticle (or particle) appearing in the p_2 direction [21].

All EPR experiments in the past 64 years have been proving the validity and completeness of QM, excluding the existence of LHV. What we add in this letter is that some times the including of antiparticle is necessary. Actually, the correct answer to the original version of EPR paradox is a unique solution. There is no other alternative.

(b) The cognition of Eqs. (6)-(9) is in total conformity with the long investigation on the problem of C, P and T reversal since the discovery of parity violation in 1956. In other words, the CPT theorem already exhibits itself as a basic postulate. The transformation of a particle to its antiparticle is not some thing which can be defined independently but a direct consequence of (newly defined) space-time reversal ($\vec{x} \rightarrow -\vec{x}, t \rightarrow -t$).

(c) The above basic symmetry should be pushed forward into Eq. (10), forming a starting point to construct the SR, the relativistic QM and the quantum field theory (QFT). ([8], [9] see also [22]).

(d) What we are discussing is the essential conformity between QM and SR. In some sense the subtle relationship between them lies in the essential equivalence of “ $+i$ ” and “ $-i$ ”. Indeed, at the very elementary level of physics, “that not forbidden is allowed” (M. Gell-Mann).

ACKNOWLEDGMENTS

We thank Profs. M-l Du, H-z Li, J-q Liang, R-k Su, B-w Xu, S-q Ying, X-y Zeng and Drs. G-h Yang, J-f Yang and Z Zhang for discussions. This work was supported in part by the NSF of China.

-
- [1] A. Einstein, B. Podolsky and N. Rosen, Phys. Rev. 47 (1935) 777-780.
 - [2] H. Guan. *Basic Concepts in Quantum Mechanics*, (High Education Press, Beijing, 1990), Chapt. 7.
 - [3] D. Bohm, *Quantum Theory* (Prentice-Hall, 1951); M. Jammer, *The Philosophy of Quantum Mechanics, the Interpretation of Quantum Mechanics in Historical Perspective*. (New York: Wiley 1974); Selleri ed. *Quantum Mechanics Versus Local Realism*.
 - [4] J. S. Bell, Physics 1 (1964) 195.
 - [5] W. Tittel, J. Brendel, B. Gisin, T. Heizog, H. Zbinden, and N. Gisin, Phys. Rev. A 57 (1998) 3229-3232.
 - [6] J. Schwinger, Proc. Nat. Acad. Sc. U.S. 44 (1958) 223.
 - [7] E. J. Konopinski and H. M. Mahmaud, Phys. Rev. 92 (1953) 1045.
 - [8] G-j Ni, Journal of Fudan University (Natural Science) No. 3-4, 125 (1974).
 - [9] G-j Ni and S-q Chen, ibid, 35 (3) 325 (1996); English version, preprint, hep-th/9508069 (1995), to appear in a book *Photon and Poincare Group* by NOVA SCIENCE PUBLISHER. Edited by V. Dvoeglazov.

- [10] T. D. Lee and C. N. Yang, Phys. Rev. 104. (1956) 254.
- [11] C. S. Wu, Ambler, Hayward, Hoppes and Hudson, Phys. Rev. 105 (1957) 1413.
- [12] T. D. Lee, R. Oehme and C. N. Yang, Phys. Rev. 106 (1957) 340.
- [13] CPLEAR Collaboration, A. Angelopoulos et al. CERN-EP/98-153, Submitted to Phys. Lett. B.
- [14] Fermi News 21 No. 21 (Oct. 30 1998), The Time Machine.
- [15] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Mechanics*, (Mc Graw-Hill Book Company 1964).
- [16] T. D. Lee and C. S. Wu, Annual Rev. Nucl., Scien. 15 (1965) 381.
- [17] G-j Ni and S-q Chen, Journal of Fudan University. (Natural Science) 36 (3) (1997) 247; G-j Ni, Preprint, hep-th/9708156, submitted to Phys. Lett. B in Sept 1997.
- [18] G-j Ni, Weimin Zhou and Jun Yan, Comparison among Klein-Gordon equation, Dirac equation and Relativistic Stationary Schrödinger equation, preprint, quant-ph/ 9901018, submitted to Phys. Rev. D.
- [19] M. Weitz, A. Huber, F. Schmidt-Kaler, D. Leibfried and T. W. Hänsch, Phys. Rev. Lett. 72, 328 (1994).
- [20] G-j Ni and Jun Yan, Lamb shift calculated by simple noncovariant method, preprint, quant-ph/9901016, submitted to Phys. Rev. A.
- [21] CPLEAR Collaboration, A. Apostolakis et. al. Phys. Lett. B 422 (1998) 339.
- [22] G-j Ni, Kexue (Science) 50 (1), 29 (1998), quant-ph/9803034; *ibid*, 50 (2) 38 (1998), quant-ph/9804013; *ibid*, 50 (3) 36 (1998), quant-ph/9806009; The English version of these three papers will be published in a book *Photon: Old Problems in light of New Ideas*, (NOVA SCIENCE PUBLISHERS), Edited by V. Dvoeglazov.